arXiv:1106.2279v1 [gr-qc] 12 Jun 2011

Time Evolution of Temperature and Entropy of a Gravitationally Collapsing de Sitter Schwarzschild Domain Wall

Evan Halstead
HEPCOS, Department of Physics, SUNY at Buffalo, Buffalo, NY 14260-1500

We investigate the time evolution of the temperature and entropy of a gravitationally collapsing de Sitter Schwarzschild domain wall as seen by an asymptotic observer. Recent work has completed this analysis for a Schwarzschild domain wall and for a 3+1 BTZ domain wall. There were some striking qualitative differences between the two. Specifically, the BTZ domain wall exhibited a decrease in entropy over time, implying that it will not collapse spontaneously. However, the BTZ domain wall contained both a cosmological constant and a different topology from the Schwarzschild domain wall, and we wish to isolate which of these is responsible for the qualitative differences. Hence, we will study the de Sitter Schwarzschild domain wall, as it has identical topology to the Schwarzschild domain wall yet also contains a cosmological constant. We utilize a wavefunctional approach where we couple a scalar field to the background of the collapsing domain wall and determine the spectrum of the radiation as a function of time. The spectrum is quasi-thermal, and the degree of thermality increases as the domain wall approaches the horizon. The fact that the distribution is thermal allows for the determination of the temperature as a function of time. Since the collapsing domain wall forms a black hole, we compare the late time temperature to the Hawking temperature and find very good agreement. The temperature also exhibits the proper scaling with the mass of the domain wall. Using the thermodynamic definition of entropy and the temperature-mass relation, we find the entropy as a function of time. We find that the temperature profile exhibits an interesting feature in the beginning of its collapse where the temperature increases for a short time before decreasing and approaching a constant. The entropy exhibits a similar feature in that there is a very small decrease in entropy before it starts to increase and eventually approach a constant. These features go away for smaller values of the cosmological constant. This suggests that 1) gravitational collapse is prevented in a certain parameter space, and 2) the probable cause of the qualitative differences in the domain wall collapse of the Schwarzschild versus BTZ domain walls is the cosmological constant and not the topology.

I. INTRODUCTION

The properties of static black holes are currently very well known. This work was pioneered by Bekenstein, Gibbons, and Hawking. They showed that quantum production of particle-antiparticle pairs near the horizon of a black hole could make it appear as though the black hole were radiating. It was also determined that the spectrum of this radiation was thermal and that the entropy of the black hole was proportional to its area [1–3]. The conventional process to determine the entropy uses the Bogolyubov method, which works by determining Bogolyubov coefficients that connect the initial vacuum state far outside the matter to the final vacuum state after the black hole is formed. As is evident, this method only utilizes the initial and final states of the system, and therefore there is no knowledge of the time dependence. Recently, Ref.[4–9] developed a method for determining the spectrum of radiation given off by a collapsing domain wall in a time dependent manner. These papers used a spherically symmetric collapsing domain wall for their analysis, and Ref.[10] applied it to a 3+1 BTZ domain wall. The most striking qualitative difference between the behavior of the two types of domain walls was that the entropy of the BTZ domain wall decreased over time, suggesting that the gravitational collapse would not occur spontaneously. The 3+1 BTZ metric, however, had two key differences from the Schwarzschild domain wall. The first and most obvious difference was a different topology. The second difference was that the BTZ domain wall metric contained a cosmological constant. For this reason, we decided to determine the time evolution of the entropy of a de Sitter Schwarzschild domain wall, as it has identical topology to the Schwarzschild domain wall but contains a cosmological constant. To complete the analysis of the time evolution of the temperature and entropy, we will first determine the equation of motion for the collapsing domain wall from an asymptotic observer's point of view. This equation of motion will be used in the quantization of a wavefunctional that contains a scalar field coupled to the background of the domain wall. Then, using the t=0 wavefunctional as a basis, we will determine the occupation number as a function of frequency. This in turn will allow for the determination of the thermodynamic quantity β and therefore the temperature. We then determine the late time temperature as a function of horizon radius. Using the thermodynamic definition of entropy, $dS = dQ/T$, we integrate to find entropy as a function of temperature, and since we know the temperature as a function of time we can find the entropy as a function of time. We compare our results against the standard Hawking results, and we comment on the results.

II. EQUATION OF MOTION OF THE DOMAIN WALL

It was shown in Ref.[4] that for a domain wall with a general outside metric

$$ds^2 = -f(r)dt^2 + \frac{1}{f(r)}dr^2 + r^2 d\Omega^2, \tag{1}$$

the classical equation of motion near the horizon from an asymptotic observer's point of view is

$$\dot{R} \approx \pm f\left(1 - \frac{1}{2}\frac{fR^4}{h^2}\right). \tag{2}$$

The dynamics for $R \sim R_H$ can be obtained by solving the equation $\dot{R} = \pm f$. For a de Sitter Schwarzschild metric,

$$f{=}1 - \frac{R_s}{R} - \frac{\Lambda R^2}{3} \tag{3}$$

and the equation of motion to first order in $R_0 - R_H$ is

$$R(t) \approx R_H + (R_0 - R_H)e^{-(1/R_H - \Lambda R_H)t}, \tag{4}$$

where R_H represents the black hole horizon radius. It should be noted that for a de Sitter Schwarzschild black hole, the horizon radius R_H is not $R_s = 2GM$. In fact, a de Sitter Schwarzschild black hole has two horizons. The outer horizon is called the cosmological horizon, and in order to produce the proper dynamics we must ensure that the collapsing domain wall starts within the cosmological horizon.

III. RADIATION WAVEFUNCTIONAL

First, we will consider the radiation given off by the domain wall during gravitational collapse. This will be done by coupling a scalar field to the background of the collapsing shell. We will decompose the scalar field as

$$\Phi = \sum_k a_k(t)u_k(r). \tag{5}$$

The exact form of $u_k(r)$ will not be important to us. To find the modes $a_k(t)$, we will insert the metrics given by Eqs.(**??**) and (**??**) into the action

$$S_\Phi = \int d^4x\sqrt{-g}g^{\mu\nu}\partial_\mu\Phi\partial_\nu\Phi, \tag{6}$$

where we are using $r = R(t)$ as the domain wall's position. The total action will be written as the sum

$$S = S_{in} + S_{out}, \tag{7}$$

where

$$S_{in} = 2\pi\int dt\int_0^{R(t)} drr^2\left(-\frac{(\partial_t\Phi)^2}{\dot{T}} + \dot{T}(\partial_r\Phi)^2\right) \tag{8}$$

is the metric inside the shell,

$$S_{out} = 2\pi\int dt\int_{R(t)}^\infty drr^2\left(-\frac{(\partial_t\Phi)^2}{f} + f(\partial_r\Phi)^2\right) \tag{9}$$

is the metric outside the shell, and

$$\frac{dT}{dt} = \frac{1}{A}\sqrt{Af - \frac{[A-f]\dot{R}^2}{f}}. \tag{10}$$

According to Ref.[12],

$$\dot{R} = f\sqrt{1 - \frac{fR^4}{h^2}} \tag{11}$$

where

$$h = \frac{f^{3/2}R^2}{\sqrt{f^2 - \dot{R}^2}}, \tag{12}$$

so dT/dt can be rewritten as

$$\frac{dT}{dt} = \frac{f}{A}\sqrt{1 + [A-f]\frac{R^4}{h}}. \tag{13}$$

In the region of interest, where $R \to R_H$ and therefore $f \to 0$, the kinetic term in S_{in} dominates over that in S_{out} while the gradient term in S_{in} is subdominant to that in S_{out}. This yields the approximate action

$$S \approx 2\pi\int dt\left[-\int_0^{R_H} drr^2\frac{(\partial_t\Phi)^2}{f} + \int_{R_H}^\infty drr^2 f(\partial_r\Phi)^2\right]. \tag{14}$$

Substituting the expansion for the scalar field Φ produces

$$\begin{aligned} S \; &= \int dt\left[-2\pi\int_0^{R_H} drr^2\frac{\sum_{k,k'}\dot{a}_k(t)\dot{a}_{k'}(t)u_k(r)u_{k'}(r)}{f}\right. \\ &+ \left. 2\pi\int_{R_H}^\infty drr^2 f\sum_{k,k'} a_k(t)a_{k'}(t)u'_k(r)u'_{k'}(r)\right]. \end{aligned} \tag{15}$$

Now let

$$M_{kk'} = 4\pi\int_0^{R_H} drr^2u_k(r)u_{k'}(r) \tag{16}$$

and

$$N_{kk'} = -4\pi\int_{R_H}^\infty drr^2 f(r)u'_k(r)u'_{k'}(r), \tag{17}$$

which allows us to rewrite (15) as

$$S = \sum_{k,k'}\int dt\left[-\frac{1}{2f}\dot{a}_kM_{kk'}\dot{a}_{k'} - \frac{1}{2}a_kN_{kk'}a_{k'}\right]. \tag{18}$$

The action (18) corresponds to the Hamiltonian

$$H = \frac{1}{2}f\Pi_kM^{-1}_{kk'}\Pi_{k'} + \frac{1}{2}a_kN_{kk'}a_{k'}, \tag{19}$$

where

$$\Pi = \frac{\partial L}{\partial \dot{a}}. \tag{20}$$

Since **M** and **N** are Hermitian matrices, it is possible to do a principle axis transformation to diagonalize them simultaneously. The exact form of them, however, is not important here. The Schrödinger Equation for a single mode will then be

$$\left[-\frac{1}{2m}f\frac{\partial^2}{\partial b^2} + \frac{1}{2}Kb^2\right]\psi(b,t) = i\frac{\partial\psi(b,t)}{\partial t}, \tag{21}$$

where m and K are the eigenvalues of **M** and **N**, respectively, and b is the eigenmode.

This, however, only represents the Hamiltonian of the radiation, not the total system. The Hamiltonian of the system can be written as

$$H_{total} = H_{wall} + H_{rad}. \tag{22}$$

Ref.[12] showed that

$$H_{wall}^2 = (f\Pi_R)^2 + f(2\pi\mu R^2)^2, \tag{23}$$

where $\mu = \sqrt{\Lambda R_H^2/3} - 2\pi\sigma G R_H$ and σ is the tension in the wall. As $R \to R_H$, $f \to 0$, and the first term on the right hand side will dominate (since $\Pi \sim f^{-3/2}$), yielding

$$H_{wall} \approx -f\Pi_R, \tag{24}$$

where we have chosen the negative solution because the shell is collapsing. The Hamiltonian of the radiation is

$$H_{rad} = \frac{f}{2m}\Pi_b^2 + \frac{K}{2}b^2. \tag{25}$$

The Hamiltonian of the entire system is therefore

$$H_{total} = -f\Pi_R + \frac{f}{2m}\Pi_b^2 + \frac{K}{2}b^2. \tag{26}$$

Note that the momentum operators have different indices, where $\Pi_R = -i\partial/\partial R$ is the momentum operator conjugate to the shell's position and $\Pi_b = -i\partial/\partial b$ is the momentum operator conjugate to the eigeinmode b. The wavefunction of the entire system is therefore a function of b, R, and t, which we will write as

$$\Psi = \Psi(b,R,t). \tag{27}$$

Using (26), the Functional Schrödinger equation can be written as

$$if\frac{\partial\Psi}{\partial R} - \frac{f}{2m}\frac{\partial^2\Psi}{\partial b^2} + \frac{K}{2}b^2\Psi = i\frac{\partial\Psi}{\partial t}. \tag{28}$$

Solving this requires knowing the classical equation of motion for the distance of the domain wall, which we will take as approximately

$$\dot{R} \approx -f, \tag{29}$$

where the negative sign was chosen because the shell is collapsing. Since R(t) is only a function of time, we can rewrite (28) as

$$if\frac{1}{\dot{R}}\frac{\partial\Psi}{\partial t} - \frac{f}{2m}\frac{\partial^2\Psi}{\partial b^2} + \frac{K}{2}b^2\Psi = i\frac{\partial\Psi}{\partial t}. \tag{30}$$

According to (29), however, $f/\dot{R} = -1$, therefore

$$-\frac{f}{2m}\frac{\partial^2\Psi}{\partial b^2} + \frac{K}{2}b^2\Psi = 2i\frac{\partial\Psi}{\partial t}. \tag{31}$$

Rewriting this in the standard simple harmonic oscillator form yields

$$\left[-\frac{1}{2m}\frac{\partial^2}{\partial b^2} + \frac{m}{2}\omega^2 b^2\right]\Psi(b,\eta) = i\frac{\partial\Psi(b,\eta)}{\partial\eta}, \tag{32}$$

where

$$\omega^2 = \frac{K}{mf} = \frac{\omega_0^2}{f} \tag{33}$$

and

$$\eta = \frac{1}{2}\int_0^t dt' f. \tag{34}$$

Here we have chosen $\eta(t=0) = 0$.

At early times, the initial vacuum state for the wavefunction ψ is that of a simple harmonic oscillator given by

$$\psi(b,\eta=0) = \left(\frac{m\omega_0}{\pi}\right)^{1/4} e^{-m\omega_0 b^2/2}. \tag{35}$$

At later times, the exact solution for the wavefunction is (see Ref.[13])

$$\psi(b,\eta) = e^{i\alpha(\eta)}\left(\frac{m}{\pi\rho^2}\right)^{1/4}\exp\left[i\frac{m}{2}\left(\frac{\rho_\eta}{\rho} + \frac{i}{\rho^2}\right)b^2\right], \tag{36}$$

where ρ_η denotes the derivative of $\rho(\eta)$ with respect to η, and $\rho(\eta)$ is the solution to the equation

$$\frac{\partial^2\rho}{\partial\eta^2} + \omega^2(\eta) = \frac{1}{\rho^3} \tag{37}$$

with initial conditions

$$\rho(0) = \frac{1}{\sqrt{\omega_0}} \tag{38}$$

$$\rho_\eta(0) = \frac{\partial\rho}{\partial\eta}\mid_0 = 0. \tag{39}$$

The phase α is given by

$$\alpha(\eta) = -\frac{1}{2}\int_0^\eta \frac{d\eta'}{\rho^2(\eta')}. \tag{40}$$

IV. OCCUPATION NUMBER OF THE RADIATION

Consider an observer with detectors that are designed to register particles for the scalar field ϕ at early times. At late times, the observer will interpret each mode b in terms of the simple harmonic oscillator states, with final frequency $\bar{\omega}$. The number of quanta in eigenmode b can be found by decomposing the wavefunction (36) into the simple harmonic oscillator states and evaluating the occupation number. The wavefunction ψ written in terms of the simple harmonic basis, $\{\varphi_n\}$, at $t = 0$ is given by

$$\psi(b, t) = \sum_n c_n(t)\varphi_n(b), \tag{41}$$

where

$$c_n(t) = \int db \varphi_n^*(b)\psi(b, t), \tag{42}$$

which is the overlap of a Gaussian with the simple harmonic basis functions. The occupation number at eigenfrequency $\bar{\omega}$ is given by the expectation value

$$N(t, \bar{\omega}) = \sum_n n|c_n|^2. \tag{43}$$

After substitution, we find that the occupation number in the eigenmode b is given by (see Appendix B in Ref.[14])

$$N(t, \bar{\omega}) = \frac{\bar{\omega}\rho^2}{4}\left[\left(1 - \frac{1}{\bar{\omega}\rho^2}\right)^2 + \left(\frac{2\rho_t}{f\bar{\omega}\rho}\right)^2\right]. \tag{44}$$

We will find that the plot is very similar to a typical Planck distribution. Therefore, we can find the temperature of the radiation by treating the occupation number of each eigenmode b as if it followed the Plank distribution,

$$N_P = \frac{1}{e^{\beta\bar{\omega}} - 1}. \tag{45}$$

For the total system, the plot of occupation number as a function of frequency is shown in Fig.(1). Notice how similar this is to a Planck distribution. In fact, as time goes on, the distribution becomes more and more Planck-like as the oscillations in the curve dampen out and the $\bar{\omega} = 0$ occupation number gets larger. If one plots $\ln(1 + 1/N)$ as a function of $\bar{\omega}$, the slope will yield β.

The plot of $ln(1 + 1/N)$ as a function of frequency is shown in Fig.(2). The straight lines represent best-fit lines whose slopes represent β at the chosen time. It should be noted at this point that β is in terms of scaled time η instead of the asymptotic observer time t. Therefore, we must rescale β back to the observer's time, which is achieved by

$$\beta^{(t)} = \frac{2\beta^{(\eta)}}{f}. \tag{46}$$

While in scaled time η the β actually diverges, in the asymptotic observer's time t the β actually approaches a constant, as shown in Fig.(3).

Finally, the temperature as a function of the observer's time can be found by inverting β, as shown in Fig.(4).

$$T(t) = \frac{1}{\beta^{(t)}} \tag{47}$$



FIG. 1: Occupation number of the radiation as a function of frequency. From bottom to top, the three lines are for t=13, t=14, and t=15, respectively.



FIG. 2: The best-fit slopes of these curves represent β.

Notice how, from Fig.(4), there is a short time where the temperature increases before it starts to decrease again. This is an effect which is much more prominent when the domain wall starts just inside the cosmological horizon, and it is suppressed the smaller the cosmological constant is. Notice also that the temperature approaches a constant at late times. According to [17], the Hawking temperature of a de Sitter Schwarzschild black hole is

$$T_H = \frac{1}{8\pi M} - \frac{\Lambda M}{2\pi}. \tag{48}$$



FIG. 3: β as a function of time for a collapsing de Sitter Schwarzschild domain wall.



FIG. 4: Temperature as a function of time for a collapsing de Sitter Schwarzschild domain wall.

From this we find that the ratio of the late time temperature in Fig.(4) to the accepted Hawking temperature is $T_{Schw\Lambda}/T_H = 1.02$. We will now determine how the late time temperature scales with the mass of the domain wall. The plot of the domain wall's late time temperature as a function of mass, along with the best-fit curve, is shown in Fig.(5). The equation of the best-fit curve is

$$T_{Schw\Lambda} = \frac{0.0397304}{M} - 0.0138729M. \tag{49}$$

Notice that the scaling with mass shows very good agreement with the expected result.

V. ENTROPY

The thermodynamic definition of entropy in terms of temperature is

$$S = \int \frac{dQ}{T}. \tag{50}$$



FIG. 5: Temperature as a function of mass for a de Sitter Schwarzschild domain wall.

Since changing the energy of the domain wall is the same as changing the mass, this can also be written as

$$S = \int \frac{dM}{T} \tag{51}$$

Therefore, if we can determine how the temperature of the domain wall depends on the mass, we will be able to find an expression for the entropy in terms of the temperature. And since we know the time evolution of the temperature, we can determine the time evolution of the entropy as well.

We insert Eq.(49) into Eq.(51), and after integrating we find that the entropy as a function of temperature is

$$\begin{aligned} S &= \frac{\gamma}{12\nu^3(T-\nu)}\left[\frac{3\gamma^2\nu\Lambda}{T-\nu} - 6\gamma^2\Lambda + \nu^2\left(6 - \frac{2\gamma^2\Lambda}{(T-\nu)^2}\right)\right] \\ &+ \frac{1}{2\nu^4}\left(-\gamma\nu^2 + \gamma^3\Lambda\right) ln\left(\gamma + \frac{\gamma\nu}{T-\nu}\right), \end{aligned} \tag{52}$$

where $\gamma = 0.0397304$ and $\nu = -0.0138729$. After normalizing the entropy such that the initial entropy is zero, we find that the entropy approaches a constant, as shown in Fig.(6). We also wish to point out here that, though the effect is small, the entropy does in fact decrease for a short time before increasing. The time dependent analysis of the 3+1 BTZ domain wall also exhibited a decrease with entropy over time, albeit in that case the effect was ever-present instead of temporary as it is here. This implies that the probable cause of the decreasing entropy is the cosmological constant and not the difference in topology. The decreasing entropy also means that for this particular choice of parameters, the domain wall will not collapse spontaneously, but rather would require an energy input to get it started. The explanation of this effect is quite intuitive, as the collapse is at odds with the expansion of space, and if the cosmological constant is large enough, then collapse will be prevented. We discovered that this feature is a function of the particular parameters that we chose. The amount of decrease is amplified when the domain wall starts close to the cosmological horizon,



FIG. 6: Entropy as a function of time for a de Sitter Schwarzschild domain wall.

and there is no longer a decrease in entropy when the cosmological constant is sufficiently small. Given how small the cosmological constant is by current measurements, it is unlikely that the collapse of a de Sitter Schwarzschild domain wall would be forbidden in our universe.

VI. CONCLUSION

We investigated the time evolution of the temperature and entropy of a de Sitter Schwarzschild domain wall (representing a spherically symmetric domain wall with a cosmological constant). This was done by coupling a scalar field to the background of the domain wall and evaluating the occupation number. From the occupation number we were able to determine β and therefore the temperature. We found that the temperature of the de Sitter Schwarzschild domain wall increased for a short time before then decreasing and approaching a constant. We found that this effect was more prominent when the initial position of the domain wall was close to the cosmological horizon. The effect is also suppressed when the cosmological constant is small. The late time temperature exhibited very good agreement with the Hawking temperature of a static black hole. It also produced scaling with mass that matched the literature. The entropy, however, was interesting because it exhibited qualities that were seen both in the Schwarzschild domain wall and in the 3+1 BTZ domain wall. Namely, overall the entropy increased and approached a constant, but there was also a short time in the beginning of the collapse when the entropy decreased by a small amount. Comparison with the results for the Schwarzschild and BTZ domain walls leads us to posit that the cause of the decrease in entropy was the cosmological constant and not the difference in topology.

Acknowledgments

The author would like to thank D. Stojkovic and E. Greenwood for their insight and their helpful conversations.

---

[1] S. W. Hawking, Commun. Math. Phys. **43**, 199 (1975) [Erratum-ibid. **46**, 206 (1976)].

[2] J. B. Hartle and S. W. Hawking, Phys. Rev. D **13**, 2188 (1976).

[3] G. W. Gibbons and S. W. Hawking, Phys. Rev. D **15**, 2752 (1977).

[4] T. Vachaspati, D. Stojkovic and L. M. Krauss, Phys. Rev. D **76**, 024005 (2007) [arXiv:gr-qc/0609024].

[5] T. Vachaspati and D. Stojkovic, Phys. Lett. B **663**, 107 (2008) [arXiv:gr-qc/0701096].

[6] E. Greenwood, JCAP **0906**, 032 (2009) [arXiv:0811.0816 [gr-qc]].

[7] E. Greenwood and D. Stojkovic, JHEP **0909**, 058 (2009) [arXiv:0806.0628 [gr-qc]]. Phys. Rev. **A45**, 1320 (19yy).

[8] J. E. Wang, E. Greenwood and D. Stojkovic, Phys. Rev. D **80**, 124027 (2009) [arXiv:0906.3250 [hep-th]].

[9] E. Greenwood, D. C. Dai and D. Stojkovic, Phys. Lett. B **692**, 226 (2010) [arXiv:1008.0869 [astro-ph.CO]].

[10] E. Halstead and P. Hao, arxiv. 0260995

[11] J. D. Bekenstein, Phys. Rev. D **7**, 2333 (1973).

[12] E. Greenwood, E. Halstead and P. Hao, JHEP **1002**, 044 (2010) [arXiv:0912.1860 [gr-qc]].

[13] C. M. A. Dantas, I. A. Pedrosa and B. Baseia,

[14] E. Greenwood, JCAP **1001**, 002 (2010) [arXiv:0910.0024 [gr-qc]].

[15] S. P. Kim, "Nonequilibrium self-interacting quantum fields in cosmology: The arXiv:hep-th/9809091.

[16] J. P. S. Lemos, Phys. Lett. B **353**, 46 (1995) [arXiv:gr-qc/9404041].

[17] I. Arraut, D. Batic and M. Nowakowski, Class. Quant. Grav. **26**, 125006 (2009) [arXiv:0810.5156 [gr-qc]].